arXiv:1302.1314v1 [math.NA] 6 Feb 2013

Error Estimates with Explicit Constants for Sinc Quadrature and Sinc Indefinite Integration over Infinite Intervals*

Tomoaki Okayama
Graduate School of Economics, Hitotsubashi University
`tokayama@econ.hit-u.ac.jp`

February 7, 2013

Abstract

The Sinc quadrature and the Sinc indefinite integration are approximation formulas for definite integration and indefinite integration, respectively, which can be applied on any interval by using an appropriate variable transformation. Their convergence rates have been analyzed on any interval. In addition, with the aim of verified computation, more explicit error bounds that are computable have been recently given on the finite interval. In this paper, such explicit error bounds are given in the remaining case: on the semi-infinite interval and the infinite interval.

1 Introduction

The "Sinc quadrature" is an approximation formula for the integral over the whole real line, expressed as

$$\int_{-\infty}^{\infty} F(x)\,\mathrm{d}x \approx h \sum_{k=-M}^{N} F(kh), \tag{1.1}$$

where M, N, h are selected appropriately depending on n. This approximation is also called the (truncated) "trapezoidal formula." It is well known that the formula (1.1) can achieve quite a fast, *exponential* convergence. Furthermore, its optimality is proved under a certain class of functions [4, 6]. Here, there are two important conditions to be satisfied: (i) the interval of integration is $(-\infty,\,\infty)$, and (ii) $|F(x)|$ decays exponentially as $x \to \pm\infty$. In other cases, users should employ an appropriate variable transformation $t = \psi(x)$, i.e., the given integral should be transformed as $\int_a^b f(t)\,\mathrm{d}t = \int_{-\infty}^{\infty} f(\psi(x))\psi'(x)\,\mathrm{d}x$, so that those two conditions are met.

In this regard, Stenger [5] considered the following four typical cases:

1. $(a,\,b) = (-\infty,\,\infty)$, and $|f(x)|$ decays algebraically as $x \to \pm\infty$,
2. $(a,\,b) = (0,\,\infty)$, and $|f(x)|$ decays algebraically as $x \to \infty$,
3. $(a,\,b) = (0,\,\infty)$, and $|f(x)|$ decays (already) exponentially as $x \to \infty$,
4. The interval $(a,\,b)$ is finite,

---

*This work was supported by the Grant-in-Aid for Young Scientists (B) (No. 24760060), JSPS, Japan.

and gave the concrete transformations to be employed in all cases:

$$\begin{aligned}
\psi_{\mathrm{SE1}}(t) &= \sinh t,\\
\psi_{\mathrm{SE2}}(t) &= \mathrm{e}^{t},\\
\psi_{\mathrm{SE3}}(t) &= \operatorname{arcsinh}(\mathrm{e}^{t}),\\
\psi_{\mathrm{SE4}}(t) &= \frac{b-a}{2}\tanh\left(\frac{t}{2}\right)+\frac{b+a}{2},
\end{aligned}$$

which are called the "Single-Exponential (SE) transformations." Takahasi–Mori [7] proposed the following improved transformations:

$$\begin{aligned}
\psi_{\mathrm{DE1}}(t) &= \sinh[(\pi/2)\sinh t],\\
\psi_{\mathrm{DE2}}(t) &= \mathrm{e}^{(\pi/2)\sinh t},\\
\psi_{\mathrm{DE3\dagger}}(t) &= \mathrm{e}^{t-\exp(-t)},\\
\psi_{\mathrm{DE4}}(t) &= \frac{b-a}{2}\tanh\left(\frac{\pi}{2}\sinh t\right)+\frac{b+a}{2},
\end{aligned}$$

which are called the "Double-Exponential (DE) transformations." In addition, in the case 3, another DE transformation

$$\psi_{\mathrm{DE3}}(t) = \log(1+\mathrm{e}^{(\pi/2)\sinh t})$$

was proposed [2] so that its inverse function can be explicitly written with elementary functions (whereas $\psi_{\mathrm{DE3\dagger}}(t)$ cannot).

Error analyses of the Sinc quadrature combined with $\psi_{\mathrm{SE1}}(t),\ldots,\psi_{\mathrm{SE4}}(t)$ have been given [5] in the following form:

$$|\mathrm{Error}| \leq C\,\mathrm{e}^{-\sqrt{2\pi d\mu n}},$$

and for $\psi_{\mathrm{DE1}}(t)$, $\psi_{\mathrm{DE2}}(t)$, $\psi_{\mathrm{DE3}}(t)$, $\psi_{\mathrm{DE4}}(t)$, their error analyses have been given [8] as

$$|\mathrm{Error}| \leq C\,\mathrm{e}^{-2\pi dn/\log(8dn/\mu)}, \tag{1.2}$$

and for $\psi_{\mathrm{DE3\dagger}}(t)$, also given [8] as

$$|\mathrm{Error}| \leq C\,\mathrm{e}^{-2\pi dn/\log(2\pi dn/\mu)}, \tag{1.3}$$

where μ indicates the decay rate of the integrand, and d denotes the width of the domain in which the transformed integrand is analytic, and C is a constant independent of n. In view of the inequalities above, we notice that the accuracy of the approximation can be guaranteed if the constant C is explicitly given in a computable form. In fact, the explicit form of C has been revealed in the case 4 (the interval is finite) [3], and the result was used for verified automatic integration [9].

The main objective of this study is to reveal the explicit form of C's in the remaining cases: 1–3 (the interval is not finite), which enables us to bound the errors by computable terms.

As a second objective, this paper improves the DE transformation in the case 3. Instead of $\psi_{\mathrm{DE3}}(t)$ or $\psi_{\mathrm{DE3\dagger}}(t)$,

$$\psi_{\mathrm{DE3\ddagger}}(t) = \log(1+\mathrm{e}^{\pi\sinh t})$$

is introduced in this paper, and it is shown that the error is estimated as

$$|\mathrm{Error}| \leq C\,\mathrm{e}^{-2\pi dn/\log(4dn/\mu)}, \tag{1.4}$$

with clarifying the constant C. The rate of (1.4) is better than (1.2) and (1.3). Furthermore, by using Sugihara's nonexistence theorem [6], it can be shown that $\psi_{\mathrm{DE3\ddagger}}$ is the best among possible variable transformations in the case 3 (although the point is not discussed in this paper and left for another occasion).

In addition to the "Sinc quadrature" described above, similar results can be given for the "Sinc indefinite integration" for indefinite integrals $\int_a^{\xi} f(t)\,\mathrm{d}t$, which is also examined in this study.

The remainder of this paper is organized as follows. Main results of this paper are stated in Sections 2 and 3; new error estimates for the Sinc quadrature are presented in Section 2, and for the Sinc indefinite integration in Section 3. All proofs of the presented theorems are given in Section 5. Numerical examples are shown in Section 4.

2 Error Estimates with Explicit Constants for the Sinc Quadrature

In this section, new error estimates for the Sinc quadrature are stated comparing with the existing results. Let us introduce some notation here. Let $\mathscr{D}_d$ be a strip domain defined by $\mathscr{D}_d = \{\zeta \in \mathbb{C} : |\operatorname{Im}\zeta| < d\}$ for $d > 0$. Furthermore, let $\mathscr{D}_d^- = \mathscr{D}_d \cap \{\zeta \in \mathbb{C} : \operatorname{Re}\zeta < 0\}$, $\mathscr{D}_d^+ = \mathscr{D}_d \cap \{\zeta \in \mathbb{C} : \operatorname{Re}\zeta \geq 0\}$. In all theorems in Sections 2 and 3, d is supposed to be a positive constant with $d < \pi/2$. For a variable transformation ψ, $\psi(\mathscr{D}_d)$ denotes the image of $\mathscr{D}_d$ by the map ψ, i.e., $\psi(\mathscr{D}_d) = \{z = \psi(\zeta) : \zeta \in \mathscr{D}_d\}$. Let $I_1 = (-\infty, \infty)$, $I_2 = I_3 = (0, \infty)$, and let us define the following three functions:

$$\begin{aligned}
E_1(z;\gamma) &= \frac{1}{(1+z^2)^{(\gamma+1)/2}},\\
E_2(z;\alpha,\beta) &= \frac{z^{\alpha-1}}{(1+z^2)^{(\alpha+\beta)/2}},\\
E_3(z;\alpha,\beta) &= \left(\frac{z}{1+z}\right)^{\alpha-1} \mathrm{e}^{-\beta z}.
\end{aligned}$$

We write $E_i(z;\gamma,\gamma)$ as $E_i(z;\gamma)$ for short.

2.1 Existing and New Error Estimates for the Sinc Quadrature with the SE Transformation

Existing error analyses for the Sinc quadrature with ψ_{SE1}, ψ_{SE2}, ψ_{SE3} are written in the following form (Theorems 2.1 and 2.2).

Theorem 2.1 (Stenger [5, Theorem 4.2.6]). Assume that f is analytic in $\psi_{\mathrm{SE1}}(\mathscr{D}_d)$, and there exist positive constants K, α, and β such that

$$|f(z)| \leq K|E_1(z;\alpha)| \tag{2.1}$$

for all $z \in \psi_{\mathrm{SE1}}(\mathscr{D}_d^-)$, and

$$|f(z)| \leq K|E_1(z;\beta)| \tag{2.2}$$

for all $z \in \psi_{\mathrm{SE1}}(\mathscr{D}_d^+)$. Let $\mu = \min\{\alpha,\, \beta\}$, let h be defined as

$$h = \sqrt{\frac{2\pi d}{\mu n}}, \tag{2.3}$$

and let M and N be defined as

$$\begin{cases} M = n, \quad N = \lceil \alpha n/\beta \rceil & (\text{if } \mu = \alpha), \\ N = n, \quad M = \lceil \beta n/\alpha \rceil & (\text{if } \mu = \beta). \end{cases} \tag{2.4}$$

Then there exists a constant C_1, independent of n, such that

$$\left| \int_{I_1} f(t)\,\mathrm{d}t - h \sum_{k=-M}^{N} f(\psi_{\mathrm{SE1}}(kh))\psi_{\mathrm{SE1}}'(kh) \right| \leq C_1 \,\mathrm{e}^{-\sqrt{2\pi d \mu n}}. \tag{2.5}$$

Theorem 2.2 (Stenger [5, Theorem 4.2.6]). The following is true for $i = 2$, 3. Assume that f is analytic in $\psi_{\mathrm{SE}i}(\mathscr{D}_d)$, and there exist positive constants K, α, and β such that

$$|f(z)| \leq K |E_i(z;\alpha,\beta)| \tag{2.6}$$

for all $z \in \psi_{\mathrm{SE}i}(\mathscr{D}_d)$. Let $\mu = \min\{\alpha,\, \beta\}$, let h be defined as (2.3), and let M and N be defined as (2.4). Then there exists a constant C_i, independent of n, such that

$$\left| \int_{I_i} f(t)\,\mathrm{d}t - h \sum_{k=-M}^{N} f(\psi_{\mathrm{SE}i}(kh))\psi_{\mathrm{SE}i}'(kh) \right| \leq C_i \,\mathrm{e}^{-\sqrt{2\pi d \mu n}}. \tag{2.7}$$

This paper explicitly estimates the constant C_i's in (2.5) and (2.7) as follows.

Theorem 2.3. Let the assumptions in Theorem 2.1 be fulfilled. Furthermore, let $\nu = \max\{\alpha,\, \beta\}$. Then the inequality (2.5) holds with

$$C_1 = \frac{2^{\nu+1}K}{\mu} \left\{ \frac{2}{(1-\,\mathrm{e}^{-\sqrt{2\pi d\mu}})\{\cos d\}^{\nu}} + 1 \right\}.$$

Theorem 2.4. Let the assumptions in Theorem 2.2 be fulfilled. Then the inequality (2.7) holds with

$$C_2 = \frac{2K}{\mu} \left\{ \frac{2}{(1-\,\mathrm{e}^{-\sqrt{2\pi d\mu}})\{\cos d\}^{(\alpha+\beta)/2}} + 1 \right\},$$

$$C_3 = \frac{2K}{\mu} \left\{ \frac{2^{(\alpha+\beta+1)/2} c_{\alpha,d}}{(1-\,\mathrm{e}^{-\sqrt{2\pi d\mu}})\{\cos d\}^{(\alpha+\beta)/2}} + 1 \right\},$$

where $c_{\alpha,d}$ is a constant defined by

$$c_{\alpha,d} = \begin{cases} \left(1 + \frac{1}{\cos d}\right)^{(1-\alpha)/2} & (\text{if } 0 < \alpha < 1), \\ 1 & (\text{if } 1 \leq \alpha). \end{cases} \tag{2.8}$$

2.2 Existing and New Error Estimates for the Sinc Quadrature with the DE Transformation

Existing error analyses for the Sinc quadrature with ψ_{DE1}, ψ_{DE2}, ψ_{DE3} and $\psi_{\mathrm{DE3}\dagger}$ are written in the following form (Theorems 2.5 and 2.6).

Theorem 2.5 (Tanaka et al. [8, Theorem 3.1])**.** The following is true for $i = 1, 2, 3$. Assume that f is analytic in $\psi_{\mathrm{DE}i}(\mathscr{D}_d)$, and there exist positive constants K and μ (with $\mu \leq 1$ in the case $i = 3$) such that

$$|f(z)| \leq K|E_i(z;\mu)|$$

for all $z \in \psi_{\mathrm{DE}i}(\mathscr{D}_d)$. Then there exists a constant C, independent of n, such that

$$\left|\int_{I_i} f(t)\,\mathrm{d}t - h\sum_{k=-n}^{n} f(\psi_{\mathrm{DE1}}(kh))\psi'_{\mathrm{DE1}}(kh)\right| \leq C\,\mathrm{e}^{-2\pi dn/\log(8dn/\mu)},$$

where

$$h = \frac{\log(8dn/\mu)}{n}. \tag{2.9}$$

Theorem 2.6 (Tanaka et al. [8, Theorem 3.1])**.** Assume that f is analytic in $\psi_{\mathrm{DE3}\dagger}(\mathscr{D}_d)$, and there exist positive constants K and μ with $\mu \leq 1$ such that

$$|f(z)| \leq K|E_3(z;\mu)|$$

for all $z \in \psi_{\mathrm{DE3}\dagger}(\mathscr{D}_d)$. Then there exists a constant C, independent of n, such that

$$\left|\int_{I_3} f(t)\,\mathrm{d}t - h\sum_{k=-n}^{n} f(\psi_{\mathrm{DE3}\dagger}(kh))\psi'_{\mathrm{DE3}\dagger}(kh)\right| \leq C\,\mathrm{e}^{-2\pi dn/\log(2\pi dn/\mu)},$$

where $h = \log(2\pi dn/\mu)/n$.

Remark 2.1. As for Theorem 2.5 with $i = 3$ and Theorem 2.6, although the condition '$\mu \leq 1$' is not assumed (only $\mu > 0$ is assumed) in the original paper [8], that condition is necessary to avoid the case where $|z/(1+z)|^{\mu-1} = \infty$ at $z = -1$ (see $E_3(z;\mu)$).

As for the case 1 (Theorem 2.5 with $i = 1$) and the case 2 (Theorem 2.5 with $i = 2$), this paper not only explicitly estimates the constant C's, but also generalizes the approximation formula from $\sum_{k=-n}^{n}$ to $\sum_{k=-M}^{N}$ as stated below. Here, x_γ is defined for $\gamma > 0$ by

$$x_\gamma = \begin{cases} \operatorname{arcsinh}\left(\dfrac{\sqrt{1+\sqrt{1-(2\pi\gamma)^2}}}{2\pi\gamma}\right) & (\text{if } 0 < \gamma < 1/(2\pi)), \\ \operatorname{arcsinh}(1) & (\text{if } 1/(2\pi) \leq \gamma). \end{cases}$$

Theorem 2.7. Assume that f is analytic in $\psi_{\mathrm{DE1}}(\mathscr{D}_d)$, and there exist positive constants K, α, and β such that (2.1) holds for all $z \in \psi_{\mathrm{DE1}}(\mathscr{D}_d^-)$, and (2.2) holds for all $z \in \psi_{\mathrm{DE1}}(\mathscr{D}_d^+)$. Let $\mu = \min\{\alpha, \beta\}$, let $\nu = \max\{\alpha, \beta\}$, let h be defined as (2.9), and let M and N be defined as

$$\begin{cases} M = n, \quad N = n - \lfloor \log(\beta/\alpha)/h \rfloor & (\text{if } \mu = \alpha), \\ N = n, \quad M = n - \lfloor \log(\alpha/\beta)/h \rfloor & (\text{if } \mu = \beta). \end{cases} \tag{2.10}$$

Furthermore, let n be taken sufficiently large so that $n \geq (\nu\,\mathrm{e})/(8d)$, $Mh \geq x_\alpha$, and $Nh \geq x_\beta$ hold. Then it holds that

$$\left|\int_{I_1} f(t)\,\mathrm{d}t - h\sum_{k=-M}^{N} f(\psi_{\mathrm{DE1}}(kh))\psi'_{\mathrm{DE1}}(kh)\right| \leq C_1\,\mathrm{e}^{-2\pi dn/\log(8dn/\mu)},$$

where C_1 is a constant independent of n, expressed as

$$C_1 = \frac{2^{\nu+1}K}{\mu}\left\{\frac{2}{(1-\mathrm{e}^{-\pi\mu\,\mathrm{e}/4})\{\cos(\frac{\pi}{2}\sin d)\}^{\nu}\cos d} + \mathrm{e}^{\pi\nu/4}\right\}.$$

Theorem 2.8. Assume that f is analytic in $\psi_{\mathrm{DE2}}(\mathscr{D}_d)$, and there exist positive constants K, α, and β such that (2.6) holds with $i = 2$ for all $z \in \psi_{\mathrm{DE2}}(\mathscr{D}_d)$. Let $\mu = \min\{\alpha, \beta\}$, let $\nu = \max\{\alpha, \beta\}$, let h be defined as (2.9), and let M and N be defined as (2.10). Furthermore, let n be taken sufficiently large so that $n \geq (\nu\,\mathrm{e})/(8d)$, $Mh \geq x_\alpha$, and $Nh \geq x_\beta$ hold. Then it holds that

$$\left|\int_{I_2} f(t)\,\mathrm{d}t - h\sum_{k=-M}^{N} f(\psi_{\mathrm{DE2}}(kh))\psi'_{\mathrm{DE2}}(kh)\right| \leq C_2\,\mathrm{e}^{-2\pi dn/\log(8dn/\mu)},$$

where C_2 is a constant independent of n, expressed as

$$C_2 = \frac{2K}{\mu}\left\{\frac{2}{(1-\,\mathrm{e}^{-\pi\mu\,\mathrm{e}/4})\{\cos(\frac{\pi}{2}\sin d)\}^{(\alpha+\beta)/2}\cos d} + \mathrm{e}^{\pi\nu/4}\right\}.$$

As for the case 3 (Theorem 2.5 with $i = 3$ and Theorem 2.6), this paper employs the improved variable transformation $\psi_{\mathrm{DE3}\ddagger}$ as described in the introduction. Then this paper gives the error estimates in the similar form to Theorems 2.7 and 2.8.

Theorem 2.9. Assume that f is analytic in $\psi_{\mathrm{DE3}\ddagger}(\mathscr{D}_d)$, and there exist positive constants K, β and α with $\alpha \leq 1$ such that (2.6) holds with $i = 3$ for all $z \in \psi_{\mathrm{DE3}\ddagger}(\mathscr{D}_d)$. Let $\mu = \min\{\alpha, \beta\}$, let $\nu = \max\{\alpha, \beta\}$, let h be defined as

$$h = \frac{\log(4dn/\mu)}{n}, \tag{2.11}$$

and let M and N be defined as (2.10). Furthermore, let n be taken sufficiently large so that $n \geq (\nu\,\mathrm{e})/(4d)$, $Mh \geq x_\alpha$, and $Nh \geq x_\beta$ hold. Then it holds that

$$\left|\int_{I_3} f(t)\,\mathrm{d}t - h\sum_{k=-M}^{N} f(\psi_{\mathrm{DE3}\ddagger}(kh))\psi'_{\mathrm{DE3}\ddagger}(kh)\right| \leq C_{3\ddagger}\,\mathrm{e}^{-2\pi dn/\log(4dn/\mu)},$$

where $C_{3\ddagger}$ is a constant independent of n, expressed as

$$C_{3\ddagger} = \frac{2K}{\mu}\left\{\frac{2(\tilde{c}_d)^{1-\alpha}}{(1-\,\mathrm{e}^{-\pi\mu\,\mathrm{e}/2})\{\cos(\frac{\pi}{2}\sin d)\}^{\alpha+\beta}\cos d} + \mathrm{e}^{\pi(1-\alpha+6\nu)/12}\right\},$$

and where $\tilde{c}_d$ is a constant expressed by using $c_d = 1 + \{1/\cos(\frac{\pi}{2}\sin d)\}$ as

$$\tilde{c}_d = \frac{1+\log(1+c_d)}{\log(1+c_d)}c_d. \tag{2.12}$$

3 Error Estimates with Explicit Constants for the Sinc Indefinite Integration

Sinc indefinite integration is an approximation formula for the indefinite integral [1], expressed as

$$\int_{-\infty}^{\xi} F(x)\,\mathrm{d}x \approx \sum_{k=-M}^{N} F(kh)J(k,h)(\xi), \quad \xi \in \mathbb{R}. \tag{3.1}$$

Here, the basis function $J(k,h)$ is defined by

$$J(k,h)(x) = h\left\{\frac{1}{2} + \frac{1}{\pi}\mathrm{Si}[\pi(x/h - k)]\right\},$$

where $\mathrm{Si}(x)$ is the so-called sine integral, defined by $\mathrm{Si}(x) = \int_0^x \{\sin(\sigma)/\sigma\}\,\mathrm{d}\sigma$. The approximation (3.1) can be combined with the SE transformation or the DE transformation [2] similar to the Sinc quadrature (1.1). This section is devoted to presenting the error estimates for the formulas.

3.1 New Error Estimates for the Sinc Indefinite Integration with the SE Transformation

This paper gives new error estimates for the Sinc indefinite integration with ψ_{SE1}, ψ_{SE2}, and ψ_{SE3} in the following form (Theorems 3.1 and 3.2).

Theorem 3.1. Assume that f is analytic in $\psi_{\mathrm{SE1}}(\mathscr{D}_d)$, and there exist positive constants K, α, and β such that (2.1) holds for all $z \in \psi_{\mathrm{SE1}}(\mathscr{D}_d^{-})$, and (2.2) holds for all $z \in \psi_{\mathrm{SE1}}(\mathscr{D}_d^{+})$. Let $\mu = \min\{\alpha,\, \beta\}$, let $\nu = \max\{\alpha,\, \beta\}$, let h be defined as

$$h = \sqrt{\frac{\pi d}{\mu n}}, \tag{3.2}$$

and let M and N be defined as (2.4). Then it holds that

$$\sup_{\tau \in I_1} \left| \int_{-\infty}^{\tau} f(t)\,\mathrm{d}t - \sum_{k=-M}^{N} f(\psi_{\mathrm{SE1}}(kh))\psi_{\mathrm{SE1}}'(kh) J(k,h)(\psi_{\mathrm{SE1}}^{-1}(\tau)) \right| \leq C_1 \,\mathrm{e}^{-\sqrt{\pi d \mu n}},$$

where C_1 is a constant independent of n, expressed as

$$C_1 = \frac{2^{\nu+1} K}{\mu} \left\{ \frac{1}{(1 - \mathrm{e}^{-2\sqrt{\pi d \mu}})\{\cos d\}^{\nu}} \sqrt{\frac{\pi}{d\mu}} + 1.1 \right\}.$$

Theorem 3.2. The following is true for $i = 2, 3$. Assume that f is analytic in $\psi_{\mathrm{SE}i}(\mathscr{D}_d)$, and there exist positive constants K, α, and β such that (2.6) holds for all $z \in \psi_{\mathrm{SE}i}(\mathscr{D}_d)$. Let $\mu = \min\{\alpha,\, \beta\}$, let h be defined as (3.2), and let M and N be defined as (2.4). Then it holds that

$$\sup_{\tau \in I_i} \left| \int_{0}^{\tau} f(t)\,\mathrm{d}t - \sum_{k=-M}^{N} f(\psi_{\mathrm{SE}i}(kh))\psi_{\mathrm{SE}i}'(kh) J(k,h)(\psi_{\mathrm{SE}i}^{-1}(\tau)) \right| \leq C_i \,\mathrm{e}^{-\sqrt{\pi d \mu n}},$$

where C_2 and C_3 are constants independent of n, expressed as

$$C_2 = \frac{2K}{\mu} \left\{ \frac{1}{(1 - \mathrm{e}^{-2\sqrt{\pi d \mu}})\{\cos d\}^{(\alpha+\beta)/2}} \sqrt{\frac{\pi}{d\mu}} + 1.1 \right\},$$

$$C_3 = \frac{2K}{\mu} \left\{ \frac{2^{(\alpha+\beta-1)/2} c_{\alpha,d}}{(1 - \mathrm{e}^{-2\sqrt{\pi d \mu}})\{\cos d\}^{(\alpha+\beta)/2}} \sqrt{\frac{\pi}{d\mu}} + 1.1 \right\},$$

and where $c_{\alpha,d}$ is a constant defined in (2.8).

3.2 New Error Estimates for the Sinc Indefinite Integration with the DE Transformation

This paper gives new error estimates for the Sinc indefinite integration with ψ_{DE1}, ψ_{DE2}, and $\psi_{\mathrm{DE3\ddagger}}$ in the following form (Theorems 3.3–3.5). Let $\epsilon_{d,\mu}^{\mathrm{DE}} = [\mathrm{e}^{-\pi d n/\log(4dn/\mu)} \log(4dn/\mu)]/n$ for short.

Theorem 3.3. Assume that f is analytic in $\psi_{\mathrm{DE1}}(\mathscr{D}_d)$, and there exist positive constants K, α, and β such that (2.1) holds for all $z \in \psi_{\mathrm{DE1}}(\mathscr{D}_d^{-})$, and (2.2) holds for all $z \in \psi_{\mathrm{DE1}}(\mathscr{D}_d^{+})$. Let $\mu = \min\{\alpha,\, \beta\}$, let $\nu = \max\{\alpha,\, \beta\}$, let h be defined as (2.11), and let M and N be defined as (2.10). Furthermore, let n be taken sufficiently large so that $n \geq (\nu\,\mathrm{e})/(4d)$, $Mh \geq x_\alpha$, and $Nh \geq x_\beta$ hold. Then it holds that

$$\sup_{\tau\in I_1}\left|\int_{-\infty}^{\tau} f(t)\,\mathrm{d}t - \sum_{k=-M}^{N} f(\psi_{\mathrm{DE1}}(kh))\psi'_{\mathrm{DE1}}(kh)J(k,h)(\psi_{\mathrm{DE1}}^{-1}(\tau))\right| \leq C_1\epsilon_{d,\mu}^{\mathrm{DE}},$$

where C_1 is a constant independent of n, expressed as

$$C_1 = \frac{2^{\nu+1}K}{\mu d}\left\{\frac{1}{(1-\,\mathrm{e}^{-\pi\mu\,\mathrm{e}/2})\{\cos(\frac{\pi}{2}\sin d)\}^{\nu}\cos d} + \mathrm{e}^{\pi(\alpha+\beta)/4}\right\}.$$

Theorem 3.4. Assume that f is analytic in $\psi_{\mathrm{DE2}}(\mathscr{D}_d)$, and there exist positive constants K, α, and β such that (2.6) holds with $i = 2$ for all $z \in \psi_{\mathrm{DE2}}(\mathscr{D}_d)$. Let $\mu = \min\{\alpha, \beta\}$, let $\nu = \max\{\alpha, \beta\}$, let h be defined as (2.11), and let M and N be defined as (2.10). Furthermore, let n be taken sufficiently large so that $n \geq (\nu\,\mathrm{e})/(4d)$, $Mh \geq x_\alpha$, and $Nh \geq x_\beta$ hold. Then it holds that

$$\sup_{\tau\in I_2}\left|\int_{0}^{\tau} f(t)\,\mathrm{d}t - \sum_{k=-M}^{N} f(\psi_{\mathrm{DE2}}(kh))\psi'_{\mathrm{DE2}}(kh)J(k,h)(\psi_{\mathrm{DE2}}^{-1}(\tau))\right| \leq C_2\epsilon_{d,\mu}^{\mathrm{DE}},$$

where C_2 is a constant independent of n, expressed as

$$C_2 = \frac{2K}{\mu d}\left\{\frac{1}{(1-\,\mathrm{e}^{-\pi\mu\,\mathrm{e}/2})\{\cos(\frac{\pi}{2}\sin d)\}^{(\alpha+\beta)/2}\cos d} + \mathrm{e}^{\pi(\alpha+\beta)/4}\right\}.$$

Theorem 3.5. Assume that f is analytic in $\psi_{\mathrm{DE3\ddagger}}(\mathscr{D}_d)$, and there exist positive constants K, β, and α with $\alpha \leq 1$ such that (2.6) holds with $i = 3$ for all $z \in \psi_{\mathrm{DE3\ddagger}}(\mathscr{D}_d)$. Let $\mu = \min\{\alpha, \beta\}$, let $\nu = \max\{\alpha, \beta\}$, let h be defined as $h = \log(2dn/\mu)/n$, and let M and N be defined as (2.10). Furthermore, let n be taken sufficiently large so that $n \geq (\nu\,\mathrm{e})/(2d)$, $Mh \geq x_\alpha$, and $Nh \geq x_\beta$ hold. Then it holds that

$$\sup_{\tau\in I_3}\left|\int_{0}^{\tau} f(t)\,\mathrm{d}t - \sum_{k=-M}^{N} f(\psi_{\mathrm{DE3\ddagger}}(kh))\psi'_{\mathrm{DE3\ddagger}}(kh)J(k,h)(\psi_{\mathrm{DE3\ddagger}}^{-1}(\tau))\right| \leq C_{3\ddagger}\epsilon_{d,2\mu}^{\mathrm{DE}},$$

where $C_{3\ddagger}$ is a constant independent of n, expressed as

$$C_{3\ddagger} = \frac{2K}{\mu d}\left\{\frac{(\tilde{c}_d)^{1-\alpha}}{(1-\,\mathrm{e}^{-\pi\mu\,\mathrm{e}})\{\cos(\frac{\pi}{2}\sin d)\}^{\alpha+\beta}\cos d} + \mathrm{e}^{\pi(1+5\alpha+6\beta)/12}\right\},$$

and where $\tilde{c}_d$ is a constant defined in (2.12).

4 Numerical Examples

In order to numerically confirm the results presented in Sections 2 and 3, let us consider the following three examples.

Example 4.1 (Case 1 [2])**.** Consider the function $f_1(t) = \sqrt{3}/(2\pi(t^2 + t + 1))$ and its definite/indefinite integral on I_1:

$$\int_{-\infty}^{\infty} f_1(t)\,\mathrm{d}t = 1, \tag{4.1}$$

$$\int_{-\infty}^{\tau} f_1(t)\,\mathrm{d}t = \frac{1}{2} + \frac{1}{\pi}\arctan\left\{\frac{2}{\sqrt{3}}\left(\tau + \frac{1}{2}\right)\right\}. \tag{4.2}$$

The integrand f_1 satisfies the assumptions in Theorem 2.3 with $\alpha = \beta = 1$, $d = 3/4$, and $K = \sqrt{3}\,\mathrm{e}$. In addition, f_1 satisfies the assumptions in Theorem 2.7 with $\alpha = \beta = 1$, $d = \pi/7$, and $K = 8\sqrt{3}/\mathrm{e}$.

Example 4.2 (Case 2 [2])**.** Consider the function $f_2(t) = 2/(\pi(1+t^2))$ and its definite/indefinite integral on I_2:

$$\int_0^{\infty} f_2(t)\,\mathrm{d}t = 1, \tag{4.3}$$

$$\int_0^{\tau} f_2(t)\,\mathrm{d}t = \frac{2}{\pi}\arctan(\tau). \tag{4.4}$$

The integrand f_2 satisfies the assumptions in Theorem 2.4 ($i = 2$) with $\alpha = \beta = 1$, $d = \cosh(1)$, and $K = 2/\pi$. In addition, f_2 satisfies the assumptions in Theorem 2.8 with $\alpha = \beta = 1$, $d = 3/2$, and $K = 2/\pi$.

Example 4.3 (Case 3 [7])**.** Consider the function $f_3(t) = \mathrm{e}^{-(1+t)}/(1+t)$ and its definite/indefinite integral on I_3:

$$\int_0^{\infty} f_3(t)\,\mathrm{d}t = \mathrm{E}_1(1), \tag{4.5}$$

$$\int_0^{\tau} f_3(t)\,\mathrm{d}t = \mathrm{E}_1(1) - \Gamma(0, 1+\tau), \tag{4.6}$$

where $\mathrm{E}_1(x)$ is the exponential integral, and $\Gamma(s, x)$ is the incomplete gamma function. The integrand f_3 satisfies the assumptions in Theorem 2.4 ($i = 3$) with $\alpha = \beta = 1$, $d = 3/2$, and $K = \mathrm{e}^{-1}$. In addition, f_3 satisfies the assumptions in Theorem 2.9 with $\alpha = \beta = 1$, $d = \log(\pi)$, and $K = \mathrm{e}$.

The numerical results are shown in Figures 1–6. All programs were written in C++ with double-precision floating-point arithmetic, and the GNU Scientific Library was used for computing special functions. In Figure 2, 'maximum error' denotes the maximum value among the errors investigated on the following 403 points: $\tau = 0$, $\pm 2^{-100}$, $\pm 2^{-99}$, $\ldots$, $\pm 2^{-1}$, $\pm 2^{0}$, $\pm 2^{1}$, $\ldots$, $\pm 2^{100}$. Similarly, in Figures 4 and 6, the errors were investigated on the 201 points (just the positive points of above), and its maximum is shown on the graphs. In each graph, we can see that the error estimate by the presented theorem (dotted line) surely bounds the actual error (solid line).

5 Proofs

5.1 In the Case of the SE Transformation

Let us have a look at the sketch of the proof first, taking the Sinc quadrature as an example. Let $F(x) = \psi_{\mathrm{SE}i}(x)\psi_{\mathrm{SE}i}'(x)$ (recall that we employ the SE transformation $t = \psi_{\mathrm{SE}i}(x)$). Then we have to evaluate the following term:

$$\left| \int_{I_i} f(t)\,\mathrm{d}t - h \sum_{k=-M}^{N} f(\psi_{\mathrm{SE}i}(kh))\psi_{\mathrm{SE}i}'(kh) \right| = \left| \int_{-\infty}^{\infty} F(x)\,\mathrm{d}x - h \sum_{k=-M}^{N} F(kh) \right|.$$

For the estimation, the function space defined below plays an important role.

Definition 5.1. Let L, R, α, β be positive constants, and d is a constant with $0 < d < \pi/2$. Then $\mathbf{L}_{L,R,\alpha,\beta}^{\mathrm{SE}}(\mathscr{D}_d)$ denotes a family of functions F that are analytic on $\mathscr{D}_d$, and satisfy for all



Figure 1. Error of the Sinc quadrature for (4.1) and its estimate.



Figure 2. Error of the Sinc indefinite integration for (4.2) and its estimate.



Figure 3. Error of the Sinc quadrature for (4.3) and its estimate.



Figure 4. Error of the Sinc indefinite integration for (4.4) and its estimate.



Figure 5. Error of the Sinc quadrature for (4.5) and its estimate.



Figure 6. Error of the Sinc indefinite integration for (4.6) and its estimate.

$\zeta \in \mathscr{D}_d$ and $x \in \mathbb{R}$ that

$$|F(\zeta)| \leq \frac{L}{|1+\mathrm{e}^{-2\zeta}|^{\alpha/2}|1+\mathrm{e}^{2\zeta}|^{\beta/2}}, \tag{5.1}$$

$$|F(x)| \leq \frac{R}{(1+\mathrm{e}^{-2x})^{\alpha/2}(1+\mathrm{e}^{2x})^{\beta/2}}. \tag{5.2}$$

If F belongs to this function space, the error of the Sinc quadrature is estimated as below. The proof is omitted since it is quite similar to the existing theorem for the case 4 [3, Theorem 2.6].

Theorem 5.1. Let $F \in \mathbf{L}^{\mathrm{SE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$, let $\mu = \min\{\alpha, \beta\}$, let h be defined as (2.3), and let M and N be defined as (2.4). Then it holds that

$$\left|\int_{-\infty}^{\infty} F(x)\,\mathrm{d}x - h\sum_{k=-M}^{N} F(kh)\right| \leq \frac{2}{\mu}\left[\frac{2L}{(1-\mathrm{e}^{-\sqrt{2\pi d\mu}})\{\cos d\}^{(\alpha+\beta)/2}} + R\right]\mathrm{e}^{-\sqrt{2\pi d\mu n}}.$$

This theorem states the desired error estimates with explicit constants for the Sinc quadrature (if $F \in \mathbf{L}^{\mathrm{SE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$, which is not confirmed yet).

For the Sinc indefinite integration, the next theorem holds. The proofs is also omitted since it is quite similar to the existing theorem for the case 4 [3, Theorem 2.9].

Theorem 5.2. Let $F \in \mathbf{L}^{\mathrm{SE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$, let $\mu = \min\{\alpha, \beta\}$, let h be defined as (3.2), and let M and N be defined as (2.4). Then it holds that

$$\begin{aligned}&\sup_{\xi\in\mathbb{R}}\left|\int_{-\infty}^{\xi} F(x)\,\mathrm{d}x - \sum_{k=-M}^{N} F(kh)J(k,h)(\xi)\right|\\ &\leq \frac{2}{\mu}\left[\frac{L}{(1-\mathrm{e}^{-2\sqrt{\pi d\mu}})\{\cos d\}^{(\alpha+\beta)/2}}\sqrt{\frac{\pi}{d\mu}} + 1.1R\right]\mathrm{e}^{-\sqrt{\pi d\mu n}}.\end{aligned}$$

This theorem states the desired error estimates with explicit constants for the Sinc indefinite integration.

In view of Theorems 5.1 and 5.2, our project is done by checking $F \in \mathbf{L}^{\mathrm{SE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$ in each case: 1, 2, 3. Let us check them one by one in what follows.

5.1.1 Proofs in the Case 1 (Theorems 2.3 and 3.1)

The claims of Theorems 2.3 and 3.1 follow from the next lemma.

Lemma 5.1. Let the assumptions in Theorem 2.3 or Theorem 3.1 be fulfilled. Then the function $F(\zeta) = f(\psi_{\mathrm{SE1}}(\zeta))\psi'_{\mathrm{SE1}}(\zeta)$ belongs to $\mathbf{L}^{\mathrm{SE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$ with $L = 2^{\nu}K/\{\cos d\}^{(\nu-\mu)/2}$ and $R = 2^{\nu}K$.

Proof. Consider the case $\alpha \leq \beta$ first. From the inequality (2.1), it follows that

$$|F(\zeta)| \leq K|E_1(\psi_{\mathrm{SE1}}(\zeta);\alpha)||\psi'_{\mathrm{SE1}}(\zeta)| = \frac{K}{|1+\mathrm{e}^{-2\zeta}|^{\alpha/2}|1+\mathrm{e}^{2\zeta}|^{\beta/2}}\cdot 2^{\alpha}|1+\mathrm{e}^{2\zeta}|^{(\beta-\alpha)/2}$$

for $\zeta \in \mathscr{D}_d$ with $\operatorname{Re}\zeta < 0$, and from the inequality (2.2), it follows that

$$|F(\zeta)| \leq K|E_1(\psi_{\mathrm{SE1}}(\zeta);\beta)||\psi'_{\mathrm{SE1}}(\zeta)| = \frac{K}{|1+\mathrm{e}^{-2\zeta}|^{\alpha/2}|1+\mathrm{e}^{2\zeta}|^{\beta/2}}\cdot\frac{2^{\beta}}{|1+\mathrm{e}^{-2\zeta}|^{(\beta-\alpha)/2}}$$

for $\zeta \in \mathscr{D}_d$ with $\operatorname{Re}\zeta \geq 0$. Setting $\zeta = x + \mathrm{i}\,y$ with $x < 0$, we have

$$\begin{aligned}
2^{\alpha}|1+\mathrm{e}^{2\zeta}|^{(\beta-\alpha)/2} &= 2^{\alpha}(1+\mathrm{e}^{2x})^{(\beta-\alpha)/2}\left\{1-\frac{\sin^2 y}{\cosh^2 x}\right\}^{(\beta-\alpha)/4} \\
&\leq 2^{\alpha}(1+\mathrm{e}^{0})^{(\beta-\alpha)/2}\left\{1-0\right\}^{(\beta-\alpha)/4} \\
&= 2^{(\alpha+\beta)/2} \leq 2^{\beta} \leq \frac{2^{\beta}}{\{\cos y\}^{(\beta-\alpha)/2}}.
\end{aligned}$$

Furthermore, setting $\zeta = x + \mathrm{i}\,y$ with $x \geq 0$, we have

$$\begin{aligned}
\frac{2^{\beta}}{|1+\mathrm{e}^{-2\zeta}|^{(\beta-\alpha)/2}} &= \frac{2^{\beta}}{(1+\mathrm{e}^{-2x})^{(\beta-\alpha)/2}\left\{1-\frac{\sin^2 y}{\cosh^2 x}\right\}^{(\beta-\alpha)/4}} \\
&\leq \frac{2^{\beta}}{(1+0)^{(\beta-\alpha)/2}\left\{1-\frac{\sin^2 y}{\cosh^2 0}\right\}^{(\beta-\alpha)/4}} = \frac{2^{\beta}}{\{\cos y\}^{(\beta-\alpha)/2}}.
\end{aligned}$$

Thus, since $\mu = \alpha$ and $\nu = \beta$ in this case, it holds for all $\zeta \in \mathscr{D}_d$ that

$$|F(\zeta)| \leq \frac{K}{|1+\mathrm{e}^{-2\zeta}|^{\alpha/2}|1+\mathrm{e}^{2\zeta}|^{\beta/2}} \cdot \frac{2^{\nu}}{\{\cos d\}^{(\nu-\mu)/2}},$$

and it holds for all $x \in \mathbb{R}$ that

$$|F(x)| \leq \frac{K}{(1+\mathrm{e}^{-2x})^{\alpha/2}(1+\mathrm{e}^{2x})^{\beta/2}} \cdot \frac{2^{\nu}}{\{\cos 0\}^{(\nu-\mu)/2}}.$$

In the case $\alpha > \beta$, the same inequalities hold. This completes the proof. ■

5.1.2 Proofs in the Case 2 (Theorems 2.4 and 3.2 with $i = 2$)

The claims of Theorems 2.4 and 3.2 ($i = 2$) follow from the next lemma.

Lemma 5.2. Let the assumptions in Theorem 2.4 or Theorem 3.2 be fulfilled with $i = 2$. Then the function $F(\zeta) = f(\psi_{\mathrm{SE2}}(\zeta))\psi_{\mathrm{SE2}}'(\zeta)$ belongs to $\mathbf{L}^{\mathrm{SE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$ with $L = K$ and $R = K$.

Proof. From the inequality (2.6) with $i = 2$, immediately (5.1) and (5.2) hold with $L = R = K$. ■

5.1.3 Proofs in the Case 3 (Theorems 2.4 and 3.2 with $i = 3$)

The claims of Theorems 2.4 and 3.2 ($i = 3$) follow from the next lemma.

Lemma 5.3. Let the assumptions in Theorem 2.4 or Theorem 3.2 be fulfilled with $i = 3$. Then the function $F(\zeta) = f(\psi_{\mathrm{SE3}}(\zeta))\psi_{\mathrm{SE3}}'(\zeta)$ belongs to $\mathbf{L}^{\mathrm{SE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$ with $L = 2^{(\alpha+\beta-1)/2}c_{\alpha,d}K$ and $R = K$, where $c_{\alpha,d}$ is a constant defined in (2.8).

For the proof, let us prepare some useful inequalities (Lemmas 5.4–5.7).

Lemma 5.4. For all $\zeta \in \overline{\mathscr{D}_{\pi/2}}$, it holds that

$$\frac{1}{\sqrt{2}}\left|\frac{\mathrm{e}^{\zeta}}{1+\mathrm{e}^{\zeta}}\right| \leq \left|\frac{\operatorname{arcsinh}(\mathrm{e}^{\zeta})}{1+\operatorname{arcsinh}(\mathrm{e}^{\zeta})}\right| \leq \sqrt{2}\left|\frac{\mathrm{e}^{\zeta}}{1+\mathrm{e}^{\zeta}}\right|. \tag{5.3}$$

Furthermore, for all $x \in \mathbb{R}$, it holds that

$$\frac{\operatorname{arcsinh}(\mathrm{e}^{x})}{1+\operatorname{arcsinh}(\mathrm{e}^{x})} \leq \frac{\mathrm{e}^{x}}{1+\mathrm{e}^{x}}. \tag{5.4}$$

Proof. Consider (5.4) first, which is proved by showing

$$\frac{t}{1+t} \leq \frac{\sinh t}{1+\sinh t}$$

for $t \geq 0$ (put $x = \log(\sinh t)$). Set $p(t) = t/(1+t)$. Since $p(t)$ is monotonically increasing, we have the desired inequality: $p(t) \leq p(\sinh t)$. Hence (5.4) is proved.

Next, consider (5.3), which is proved by showing $|g(\zeta)| \leq \sqrt{2}$ and $|1/g(\zeta)| \leq \sqrt{2}$, where

$$g(\zeta) = \frac{\operatorname{arcsinh}(\mathrm{e}^{\zeta})}{1+\operatorname{arcsinh}(\mathrm{e}^{\zeta})}\frac{1+\mathrm{e}^{\zeta}}{\mathrm{e}^{\zeta}}.$$

The functions g and $1/g$ are analytic on $\mathscr{D}_{\pi/2}$ (and continuous on $\overline{\mathscr{D}_{\pi/2}}$). Therefore, by the maximum modulus principle, $|g(\zeta)|$ and $|1/g(\zeta)|$ have their maximum on the boundary of $\mathscr{D}_{\pi/2}$, i.e., $|\operatorname{Im}\zeta| = \pi/2$. It is sufficient to consider $z = x + \mathrm{i}\,(\pi/2)$ for $x \in \mathbb{R}$.

Let us show $1/|g(z)| \leq \sqrt{2}$ first, which is relatively easy. Setting $X = \log[|1 - \mathrm{e}^{2x}| + \mathrm{e}^{2x}]$ and $Y = \arg[\sqrt{1-\mathrm{e}^{2x}} + \mathrm{i}\,\mathrm{e}^{x}]$, we have

$$|g(z)|^2 = |g(x+\mathrm{i}\,\pi/2)|^2 = (1+\mathrm{e}^{-2x})\frac{X^2+Y^2}{(X+1)^2+Y^2}.$$

Furthermore, from

$$X = \begin{cases} \log[2\,\mathrm{e}^{2x}-1] & (\text{if } x \geq 0), \\ 0 & (\text{if } x < 0), \end{cases}$$

$$Y = \begin{cases} \pi/2 & (\text{if } x \geq 0), \\ \arctan(\mathrm{e}^{x}/\sqrt{1-\mathrm{e}^{2x}}) & (\text{if } x < 0), \end{cases}$$

it holds for $x \geq 0$ that

$$\begin{aligned} \frac{1}{|g(z)|^2} &= \frac{1}{1+\mathrm{e}^{-2x}} \cdot \frac{(1+\log[2\,\mathrm{e}^{2x}-1])^2 + (\pi/2)^2}{\log^2[2\,\mathrm{e}^{2x}-1] + (\pi/2)^2} \\ &\leq \frac{1}{1+\mathrm{e}^{-2x}} \cdot \frac{(1+\{\sqrt{1+\pi^2}-1\}/2)^2 + (\pi/2)^2}{(\{\sqrt{1+\pi^2}-1\}/2)^2 + (\pi/2)^2} < 2, \end{aligned}$$

and it holds for $x < 0$ that

$$\frac{1}{|g(z)|^2} = \frac{1}{1+\mathrm{e}^{-2x}}\left\{1 + \frac{1}{\arctan^2[\mathrm{e}^{x}/\sqrt{1-\mathrm{e}^{2x}}]}\right\} \leq \frac{1}{1+\mathrm{e}^{-2x}}\left\{1+\frac{1}{(\mathrm{e}^{x})^2}\right\} = 1.$$

Thus $|1/g(x+\mathrm{i}\,\pi/2)|^2 \leq 2$ holds for all $x \in \mathbb{R}$.

Next, let us show $|g(z)| \leq \sqrt{2}$. It holds for $x \geq 0$ that

$$|g(z)|^2 = (1+\mathrm{e}^{-2x})\frac{\log^2[2\,\mathrm{e}^{2x}-1] + (\pi/2)^2}{(1+\log[2\,\mathrm{e}^{2x}-1])^2 + (\pi/2)^2} \leq (1+1)\cdot 1 = 2.$$

For $x < 0$, we have

$$|g(z)|^2 = (1+\mathrm{e}^{-2x})\frac{\arctan^2[\mathrm{e}^{x}/\sqrt{1-\mathrm{e}^{2x}}]}{1+\arctan^2[\mathrm{e}^{x}/\sqrt{1-\mathrm{e}^{2x}}]},$$

and let us bound the right hand side in the following three cases:

a) $-\frac{1}{2}\log[1+\frac{8}{\pi^2}] \leq x < 0$,

b) $-\frac{1}{2}\log[1+\frac{18}{\pi^2}] \leq x \leq -\frac{1}{2}\log[\frac{4}{3}]$ (note that $-\frac{1}{2}\log[1+\frac{8}{\pi^2}] < -\frac{1}{2}\log[\frac{4}{3}]$),

c) $x < -\frac{1}{2}\log[1+\frac{18}{\pi^2}]$.

In the case of a), it holds that

$$|g(z)|^2 \leq (1+\mathrm{e}^{-2x})\frac{(\pi/2)^2}{1+(\pi/2)^2} \leq \left(1+\left(1+\frac{8}{\pi^2}\right)\right)\frac{(\pi/2)^2}{1+(\pi/2)^2} = 2.$$

In the case of b), it holds that

$$|g(z)|^2 \leq (1+\mathrm{e}^{-2x})\frac{(\pi/3)^2}{1+(\pi/3)^2} \leq \left(1+\left(1+\frac{18}{\pi^2}\right)\right)\frac{(\pi/3)^2}{1+(\pi/3)^2} = 2.$$

In the case of c), it holds that

$$\begin{aligned}|g(z)|^2 &\leq (1+\mathrm{e}^{-2x})\frac{[\mathrm{e}^{x}/\sqrt{1-\mathrm{e}^{2x}}]^2}{1+[\mathrm{e}^{x}/\sqrt{1-0}]^2}\\ &= \frac{1}{2}\left(1-\frac{1}{\tanh x}\right) \leq \frac{1}{2}\left(1-\frac{1}{\tanh(-\frac{1}{2}\log[1+\frac{18}{\pi^2}])}\right) = 1+\frac{\pi^2}{18} < 2.\end{aligned}$$

Thus $|g(x+\mathrm{i}\,\pi/2)|^2 \leq 2$ for all $x \in \mathbb{R}$. This completes the proof. ∎

Lemma 5.5. For all $\zeta \in \overline{\mathscr{D}_{\pi/2}}$ and $x \in \mathbb{R}$, we have

$$\frac{1}{|\,\mathrm{e}^{\zeta}+\sqrt{1+\mathrm{e}^{2\zeta}}|} \leq \frac{\sqrt{2}}{|1+\mathrm{e}^{\zeta}|}, \tag{5.5}$$

$$\frac{1}{\mathrm{e}^{x}+\sqrt{1+\mathrm{e}^{2x}}} \leq \frac{1}{1+\mathrm{e}^{x}}. \tag{5.6}$$

Proof. Consider (5.6) first, which is proved by showing

$$g(x) = \frac{1+\mathrm{e}^{x}}{\mathrm{e}^{x}+\sqrt{1+\mathrm{e}^{2x}}} \leq 1$$

for $x \in \mathbb{R}$. Since

$$g'(x) = -\,\mathrm{e}^{x}\left(1-\frac{\mathrm{e}^{x}}{\sqrt{1+\mathrm{e}^{2x}}}\right)\left((1+\mathrm{e}^{x})-\sqrt{1+\mathrm{e}^{2x}}\right) \leq 0,$$

we have $g(x) \leq \lim_{x\to-\infty}\{g(x)\} = 1$, which is the desired result.

Next, consider (5.5), which is proved by showing $|g(\zeta)| \leq \sqrt{2}$. The function g is analytic on $\mathscr{D}_{\pi/2}$, (and continuous on $\overline{\mathscr{D}_{\pi/2}}$). Therefore, by the maximum modulus principle, $|g(\zeta)|$ has its maximum on the boundary of $\mathscr{D}_{\pi/2}$, i.e., $|\operatorname{Im}\zeta| = \pi/2$. It is sufficient to consider $z = x+\mathrm{i}\,(\pi/2)$ for $x \in \mathbb{R}$. In the case $x \geq 0$, we have

$$|g(z)|^2 = \left|\frac{1}{1+\sqrt{1-\mathrm{e}^{-2x}}} - \mathrm{i}\,\frac{\mathrm{e}^{-x}}{1+\sqrt{1-\mathrm{e}^{-2x}}}\right|^2 = \frac{1+\mathrm{e}^{-2x}}{(1+\sqrt{1-\mathrm{e}^{-2x}})^2} \leq 1+\mathrm{e}^{-2x} \leq 2.$$

In the case $x < 0$, we have

$$|g(z)|^2 = \left|\sqrt{1-\mathrm{e}^{2x}}+\mathrm{e}^{2x}-\mathrm{i}\,(1-\sqrt{1-\mathrm{e}^{2x}})\,\mathrm{e}^{x}\right|^2 = 1+\mathrm{e}^{2x} \leq 2.$$

This completes the proof. ∎

Lemma 5.6. For all $\zeta \in \mathscr{D}_{\pi/2}$, it holds that

$$\frac{1}{|1+\mathrm{e}^{\zeta}|} \leq \frac{1}{|1+\mathrm{e}^{2\zeta}|^{1/2}}.$$

Proof. Let x, $y \in \mathbb{R}$ with $|y| < \pi/2$, and let $\zeta = x + \mathrm{i}\,y$. Then we have

$$\left|\frac{1}{1+\mathrm{e}^{2\zeta}}\right| - \left|\frac{1}{1+\mathrm{e}^{\zeta}}\right|^2 = \frac{\mathrm{e}^{-x}}{2(\cosh(x)+\cos(y))}\sqrt{\frac{1+2\cosh(x)\cos(y)}{\cosh^2(x)-\sin^2(y)}} \geq 0,$$

which completes the proof. ■

Lemma 5.7. Let x, $y \in \mathbb{R}$ with $|y| < \pi/2$. Then it holds that

$$\sup_{x\in\mathbb{R}}\left|\frac{\cosh^2[(x+\mathrm{i}\,y)/2]}{\cosh(x+\mathrm{i}\,y)}\right| \leq \frac{\cos^2(y/2)}{\cos y} = \frac{1}{2}\left(1+\frac{1}{\cos y}\right).$$

Proof. Firstly, we have

$$\left|\frac{\cosh^2[(x+\mathrm{i}\,y)/2]}{\cosh(x+\mathrm{i}\,y)}\right| = \frac{\cosh(x)+\cos(y)}{2\sqrt{\cosh^2(x)-\sin^2(y)}} \leq \frac{\cosh(x)+\cos(y)}{2\sqrt{\cosh^2(x)\{1-\sin^2(y)\}}}$$

which then is estimated as

$$\frac{\cosh(x)+\cos(y)}{2\sqrt{\cosh^2(x)\{1-\sin^2(y)\}}} = \frac{\cosh(x)+\cos(y)}{2\cosh(x)\cos(y)} \leq \frac{1+\cos(y)}{2\cdot 1\cdot\cos(y)}.$$

This completes the proof. ■

Then Lemma 5.3 is proved as follows.

Proof. From the inequality (2.6) with $i = 3$, it follows that

$$|F(\zeta)| \leq K\left|\frac{\operatorname{arcsinh}(\mathrm{e}^{\zeta})}{1+\operatorname{arcsinh}(\mathrm{e}^{\zeta})}\right|^{\alpha-1}\left|\frac{1}{\mathrm{e}^{\zeta}+\sqrt{1+\mathrm{e}^{2\zeta}}}\right|^{\beta}\left|\frac{1}{1+\mathrm{e}^{-2\zeta}}\right|^{1/2}.$$

Consider the case $1 \leq \alpha$ first. From Lemmas 5.4, 5.5, and 5.6, it holds that

$$\begin{aligned}|F(\zeta)| &\leq K\left|\frac{\sqrt{2}}{1+\mathrm{e}^{-\zeta}}\right|^{\alpha-1}\left|\frac{\sqrt{2}}{1+\mathrm{e}^{\zeta}}\right|^{\beta}\left|\frac{1}{1+\mathrm{e}^{-2\zeta}}\right|^{1/2}\\ &\leq 2^{(\alpha+\beta-1)/2}K\left|\frac{1}{1+\mathrm{e}^{-2\zeta}}\right|^{(\alpha-1)/2}\left|\frac{1}{1+\mathrm{e}^{2\zeta}}\right|^{\beta/2}\left|\frac{1}{1+\mathrm{e}^{-2\zeta}}\right|^{1/2}\end{aligned}$$

for all $\zeta \in \mathscr{D}_d$. For $x \in \mathbb{R}$, it holds that

$$\begin{aligned}|F(x)| &\leq K\left(\frac{1}{1+\mathrm{e}^{-x}}\right)^{\alpha-1}\left(\frac{1}{1+\mathrm{e}^{x}}\right)^{\beta}\left(\frac{1}{1+\mathrm{e}^{-2x}}\right)^{1/2}\\ &\leq K\left(\frac{1}{1+\mathrm{e}^{-2x}}\right)^{(\alpha-1)/2}\left(\frac{1}{1+\mathrm{e}^{2x}}\right)^{\beta/2}\left(\frac{1}{1+\mathrm{e}^{-2x}}\right)^{1/2}.\end{aligned}$$

This completes the proof for $1 \leq \alpha$.

Next, consider the case $0 < \alpha < 1$. From Lemmas 5.4, 5.5, and 5.6, it holds that

$$\begin{aligned}|F(\zeta)| &\leq K \left|\frac{1+\mathrm{e}^{-\zeta}}{\sqrt{2}}\right|^{1-\alpha} \left|\frac{\sqrt{2}}{1+\mathrm{e}^{\zeta}}\right|^{\beta} \left|\frac{1}{1+\mathrm{e}^{-2\zeta}}\right|^{1/2} \\ &= K \left|\frac{(1+\mathrm{e}^{-\zeta})^2}{2(1+\mathrm{e}^{-2\zeta})}\right|^{(1-\alpha)/2} \left|\frac{\sqrt{2}}{1+\mathrm{e}^{\zeta}}\right|^{\beta} \left|\frac{1}{1+\mathrm{e}^{-2\zeta}}\right|^{\alpha/2}.\end{aligned}$$

Furthermore, from Lemma 5.7, it holds for $\zeta = x + \mathrm{i}\, y \in \mathscr{D}_d$ that

$$\left|\frac{(1+\mathrm{e}^{-\zeta})^2}{2(1+\mathrm{e}^{-2\zeta})}\right| = \left|\frac{\cosh^2(\zeta/2)}{\cosh(\zeta)}\right| \leq \frac{1}{2}\left(1 + \frac{1}{\cos y}\right).$$

Then using Lemma 5.6 we have

$$|F(\zeta)| \leq 2^{(\alpha+\beta-1)} K \left(1 + \frac{1}{\cos d}\right)^{(1-\alpha)/2} \left|\frac{1}{1+\mathrm{e}^{2\zeta}}\right|^{\beta/2} \left|\frac{1}{1+\mathrm{e}^{-2\zeta}}\right|^{\alpha/2}$$

for all $\zeta \in \mathscr{D}_d$. For $x \in \mathbb{R}$, it holds that

$$\begin{aligned}|F(x)| &\leq K \left(\frac{1+\mathrm{e}^{-x}}{\sqrt{2}}\right)^{1-\alpha} \left(\frac{1}{1+\mathrm{e}^{x}}\right)^{\beta} \left(\frac{1}{1+\mathrm{e}^{-2x}}\right)^{1/2} \\ &= K \left(\frac{(1+\mathrm{e}^{-x})^2}{2(1+\mathrm{e}^{-2x})}\right)^{(1-\alpha)/2} \left(\frac{1}{1+\mathrm{e}^{x}}\right)^{\beta} \left(\frac{1}{1+\mathrm{e}^{-2x}}\right)^{\alpha/2} \\ &\leq K \left(\frac{1}{2}\left(1+\frac{1}{\cos 0}\right)\right)^{(1-\alpha)/2} \left(\frac{1}{1+\mathrm{e}^{2x}}\right)^{\beta/2} \left(\frac{1}{1+\mathrm{e}^{-2x}}\right)^{\alpha/2}.\end{aligned}$$

This completes the proof for $0 < \alpha < 1$. ■

5.2 In the Case of the DE Transformation

We need the following definition in the case of the DE transformation.

Definition 5.2. Let L, R, α, β be positive constants, and d is a constant with $0 < d < \pi/2$. Then $\mathbf{L}^{\mathrm{DE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$ denotes a family of functions F that are analytic on $\mathscr{D}_d$, and satisfy for all $\zeta \in \mathscr{D}_d$ and $x \in \mathbb{R}$ that

$$|F(\zeta)| \leq \frac{(\pi/2)L|\cosh\zeta|}{|1+\mathrm{e}^{-\pi\sinh\zeta}|^{\alpha/2}|1+\mathrm{e}^{\pi\sinh\zeta}|^{\beta/2}}, \tag{5.7}$$

$$|F(x)| \leq \frac{(\pi/2)R\cosh x}{(1+\mathrm{e}^{-\pi\sinh x})^{\alpha/2}(1+\mathrm{e}^{\pi\sinh x})^{\beta/2}}. \tag{5.8}$$

If F belongs to this function space, the errors of the Sinc quadrature and the Sinc indefinite integration are estimated as below. The proofs are omitted since they are quite similar to the existing theorems for the case 4 [3, Theorems 2.14 and 2.16].

Theorem 5.3. Let $F \in \mathbf{L}^{\mathrm{DE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$, let $\mu = \min\{\alpha, \beta\}$, let $\nu = \max\{\alpha, \beta\}$, let h be defined as (2.9), and let M and N be defined as (2.10). Furthermore, let n be taken sufficiently large so that $n \geq (\nu\,\mathrm{e})/(8d)$, $Mh \geq x_\alpha$, and $Nh \geq x_\beta$ hold. Then,

$$\begin{aligned}&\left|\int_{-\infty}^{\infty} F(x)\,\mathrm{d}x - h\sum_{k=-M}^{N} F(kh)\right| \\ &\leq \frac{2}{\mu}\left[\frac{2L}{(1-\mathrm{e}^{-\pi\mu\,\mathrm{e}/4})\{\cos(\frac{\pi}{2}\cos d)\}^{(\alpha+\beta)/2}\cos d} + R\,\mathrm{e}^{\pi\nu/4}\right] \mathrm{e}^{-2\pi d n/\log(8dn/\mu)}.\end{aligned}$$

Theorem 5.4. Let $F \in \mathbf{L}^{\mathrm{DE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$, let $\mu = \min\{\alpha,\, \beta\}$, let $\nu = \max\{\alpha,\, \beta\}$, let h be defined as (2.11), and let M and N be defined as (2.10). Furthermore, let n be taken sufficiently large so that $n \geq (\nu\,\mathrm{e})/(4d)$, $Mh \geq x_\alpha$, and $Nh \geq x_\beta$ hold. Then,

$$\begin{aligned}
&\sup_{\xi\in\mathbb{R}}\left|\int_{-\infty}^{\xi} F(x)\,\mathrm{d}x - \sum_{k=-M}^{N} F(kh)J(k,h)(\xi)\right| \\
&\leq \frac{2}{\mu d}\left[\frac{L}{(1-\,\mathrm{e}^{-\pi\mu\,\mathrm{e}/2})\{\cos(\frac{\pi}{2}\sin d)\}^{(\alpha+\beta)/2}\cos d} + R\,\mathrm{e}^{\pi(\alpha+\beta)/4}\right]\epsilon^{\mathrm{DE}}_{d,\mu}.
\end{aligned}$$

In view of Theorems 5.3 and 5.4, our project is done by checking $F \in \mathbf{L}^{\mathrm{DE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$ in each case: 1, 2, 3. Let us check them one by one in what follows. The next lemma is useful for the proofs.

Lemma 5.8 (Okayama et al. [3, Lemma 4.22])**.** Let $x,\, y \in \mathbb{R}$ with $|y| < \pi/2$, and let $\zeta = x + \mathrm{i}\,y$. Then,

$$\begin{aligned}
\left|\frac{1}{1+\,\mathrm{e}^{\pi\sinh\zeta}}\right| &\leq \frac{1}{(1+\,\mathrm{e}^{\pi\sinh(x)\cos y})\cos(\frac{\pi}{2}\sin y)},\\
\left|\frac{1}{1+\,\mathrm{e}^{-\pi\sinh\zeta}}\right| &\leq \frac{1}{(1+\,\mathrm{e}^{-\pi\sinh(x)\cos y})\cos(\frac{\pi}{2}\sin y)}.
\end{aligned}$$

5.2.1 Proofs in the Case 1 (Theorems 2.7 and 3.3)

The claims of Theorems 2.7 and 3.3 follow from the next lemma.

Lemma 5.9. Let the assumptions in Theorem 2.7 or Theorem 3.3 be fulfilled. Then the function $F(\zeta) = f(\psi_{\mathrm{DE1}}(\zeta))\psi'_{\mathrm{DE1}}(\zeta)$ belongs to $\mathbf{L}^{\mathrm{DE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$ with $L = 2^\nu K/\{\cos(\frac{\pi}{2}\sin d)\}^{(\nu-\mu)/2}$ and $R = 2^\nu K$.

Proof. Consider the case $\alpha \leq \beta$ first. From the inequality (2.1), it follows that

$$|F(\zeta)| \leq \frac{K}{|1+\,\mathrm{e}^{-\pi\sinh\zeta}|^{\alpha/2}|1+\,\mathrm{e}^{\pi\sinh\zeta}|^{\beta/2}}\cdot 2^\alpha|1+\,\mathrm{e}^{\pi\sinh\zeta}|^{(\beta-\alpha)/2}$$

for $\zeta \in \mathscr{D}_d$ with $\operatorname{Re}\zeta < 0$, and from the inequality (2.2), it follows that

$$|F(\zeta)| \leq \frac{K}{|1+\,\mathrm{e}^{-\pi\sinh\zeta}|^{\alpha/2}|1+\,\mathrm{e}^{\pi\sinh\zeta}|^{\beta/2}}\cdot\frac{2^\beta}{|1+\,\mathrm{e}^{-\pi\sinh\zeta}|^{(\beta-\alpha)/2}}$$

for $\zeta \in \mathscr{D}_d$ with $\operatorname{Re}\zeta \geq 0$. Setting $\zeta = x + \mathrm{i}\,y$ with $x < 0$, we have

$$\begin{aligned}
&2^\alpha|1+\,\mathrm{e}^{\pi\sinh\zeta}|^{(\beta-\alpha)/2}\\
&= 2^\alpha(1+\,\mathrm{e}^{\pi\sinh(x)\cos y})^{(\beta-\alpha)/2}\left\{1-\frac{\sin^2(\pi\cosh(x)\sin y)}{\cosh^2(\pi\sinh(x)\cos y)}\right\}^{(\beta-\alpha)/4}\\
&\leq 2^\alpha(1+\,\mathrm{e}^0)^{(\beta-\alpha)/2}\left\{1-0\right\}^{(\beta-\alpha)/4} = 2^{(\alpha+\beta)/2} \leq 2^\beta \leq \frac{2^\beta}{\{\cos(\frac{\pi}{2}\sin y)\}^{(\beta-\alpha)/2}}.
\end{aligned}$$

Furthermore, setting $\zeta = x + \mathrm{i}\,y$ with $x \geq 0$ and using Lemma 5.8, we have

$$\begin{aligned}
\frac{2^\beta}{|1+\,\mathrm{e}^{-\pi\sinh\zeta}|^{(\beta-\alpha)/2}} &\leq \frac{2^\beta}{(1+\,\mathrm{e}^{-\pi\sinh(x)\cos y})^{(\beta-\alpha)/2}\{\cos(\frac{\pi}{2}\sin y)\}^{(\beta-\alpha)/2}}\\
&\leq \frac{2^\beta}{(1+0)\{\cos(\frac{\pi}{2}\sin y)\}^{(\beta-\alpha)/2}}.
\end{aligned}$$

Thus, since $\mu = \alpha$ and $\nu = \beta$ in this case, it holds for all $\zeta \in \mathscr{D}_d$ that

$$|F(\zeta)| \leq \frac{K}{|1+\mathrm{e}^{-\pi\sinh\zeta}|^{\alpha/2}|1+\mathrm{e}^{\pi\sinh\zeta}|^{\beta/2}} \cdot \frac{2^{\nu}}{\{\cos(\frac{\pi}{2}\sin d)\}^{(\nu-\mu)/2}},$$

and it holds for all $x \in \mathbb{R}$ that

$$|F(x)| \leq \frac{K}{(1+\mathrm{e}^{-\pi\sinh x})^{\alpha/2}(1+\mathrm{e}^{\pi\sinh x})^{\beta/2}} \cdot \frac{2^{\nu}}{\{\cos(\frac{\pi}{2}\sin 0)\}^{(\nu-\mu)/2}}.$$

In the case $\alpha > \beta$, the same inequalities hold. This completes the proof. ■

5.2.2 Proofs in the Case 2 (Theorems 2.8 and 3.4)

The claims of Theorems 2.8 and 3.4 follow from the next lemma.

Lemma 5.10. Let the assumptions in Theorem 2.8 or Theorem 3.4 be fulfilled. Then the function $F(\zeta) = f(\psi_{\mathrm{DE2}}(\zeta))\psi'_{\mathrm{DE2}}(\zeta)$ belongs to $\mathbf{L}^{\mathrm{DE}}_{L,R,\alpha,\beta}(\mathscr{D}_d)$ with $L = K$ and $R = K$.

Proof. From the inequality (2.6) with $i = 2$, immediately (5.7) and (5.8) hold with $L = R = K$. ■

5.2.3 Proofs in the Case 3 (Theorems 2.9 and 3.5)

The claims of Theorems 2.9 and 3.5 follow from the next lemma.

Lemma 5.11. Let the assumptions in Theorem 2.9 or Theorem 3.5 be fulfilled. Then the function $F(\zeta) = f(\psi_{\mathrm{DE3\ddagger}}(\zeta))\psi'_{\mathrm{DE3\ddagger}}(\zeta)$ belongs to $\mathbf{L}^{\mathrm{DE}}_{L,R,2\alpha,2\beta}(\mathscr{D}_d)$ with $L = 2(\tilde{c}_d)^{1-\alpha}K$ and $R = 2(\mathrm{e}^{\pi/12})^{1-\alpha}K$, where $\tilde{c}_d$ is a constant defined in (2.12).

For the proof, we need the next estimate.

Lemma 5.12. Let d be a constant with $0 < d < \pi/2$. Then,

$$\sup_{\zeta\in\mathscr{D}_d}\left|\frac{1+\log(1+\mathrm{e}^{\pi\sinh\zeta})}{\log(1+\mathrm{e}^{\pi\sinh\zeta})}\cdot\frac{1}{1+\mathrm{e}^{-\pi\sinh\zeta}}\right| \leq \tilde{c}_d, \tag{5.9}$$

$$\sup_{x\in\mathbb{R}}\left\{\frac{1+\log(1+\mathrm{e}^{\pi\sinh x})}{\log(1+\mathrm{e}^{\pi\sinh x})}\cdot\frac{1}{1+\mathrm{e}^{-\pi\sinh x}}\right\} \leq \mathrm{e}^{\pi/12}, \tag{5.10}$$

where $\tilde{c}_d$ is a constant defined in (2.12).

Proof. Consider (5.10) first, which is proved by showing

$$p(t) = \frac{1+t}{t}(1-\mathrm{e}^{-t}) \leq 1 + \frac{\log 6}{6} \quad \left(< \mathrm{e}^{\pi/12}\right)$$

for $t \geq 0$ (put $t = \log(1+\mathrm{e}^{\pi\sinh x})$). Let λ be a value with $\log 6 < \lambda < \log 7$ so that $p'(\lambda) = (1+\lambda+\lambda^2-\mathrm{e}^{\lambda})/(\mathrm{e}^{\lambda}\lambda^2) = 0$. Then clearly $p(t) \leq p(\lambda)$ holds. Furthermore, using the relation $1+\lambda+\lambda^2 = \mathrm{e}^{\lambda}$, we have

$$p(\lambda) = \frac{1+\lambda}{\lambda}\left(\frac{\mathrm{e}^{\lambda}-1}{\mathrm{e}^{\lambda}}\right) = \frac{1+\lambda}{\lambda}\left(\frac{\lambda(1+\lambda)}{\mathrm{e}^{\lambda}}\right) = \frac{(1+\lambda+\lambda^2)+\lambda}{\mathrm{e}^{\lambda}} = 1+\lambda\,\mathrm{e}^{-\lambda}.$$

Since the function $1 + x\,\mathrm{e}^{-x}$ is monotonically decreasing, $1+\lambda\,\mathrm{e}^{-\lambda} < 1 + (\log 6)\,\mathrm{e}^{-\log 6}$ holds (note that $\log 6 < \lambda$). Hence (5.10) is proved.

Next, consider (5.9). By the maximum modulus principle, that is proved by showing

$$|p(\xi)| \leq \frac{1+\log(2+\mathrm{e}^{\gamma})}{\log(2+\mathrm{e}^{\gamma})}(1+\mathrm{e}^{\gamma}) = \tilde{c}_d, \tag{5.11}$$

where $\xi = \log(1+\mathrm{e}^{\pi\sinh(x+\mathrm{i}\,d)})$ and $\gamma = -\log[\cos(\frac{\pi}{2}\sin d)]$. Here, notice that

$$\operatorname{Re}\xi = \log|1+\mathrm{e}^{\pi\sinh(x+\mathrm{i}\,d)}| \geq \log\left[(1+\mathrm{e}^{\pi\sinh(x)\cos d})\cos(\tfrac{\pi}{2}\sin d)\right] \geq -\gamma$$

holds from Lemma 5.8. Let us bound $|p(\xi)|$ in the two cases: a) $|\xi| \leq \log(2+\mathrm{e}^{\gamma})$, and b) $|\xi| > \log(2+\mathrm{e}^{\gamma})$. In the case of a), it holds that

$$|p(\xi)| = \left|(1+\xi)\sum_{k=1}^{\infty}\frac{(-\xi)^{k-1}}{k!}\right| \leq (1+|\xi|)\sum_{k=1}^{\infty}\frac{|\xi|^{k-1}}{k!} = \frac{1+|\xi|}{|\xi|}(\mathrm{e}^{|\xi|}-1).$$

Furthermore, since $(1+x)(\mathrm{e}^{x}-1)/x$ is monotonically increasing, we have (5.11). In the case of b), since $\operatorname{Re}\xi \geq \gamma$, it holds that

$$|p(\xi)| \leq \frac{1+|\xi|}{|\xi|}(1+|\mathrm{e}^{-\xi}|) = \frac{1+|\xi|}{|\xi|}(1+\mathrm{e}^{-\operatorname{Re}\xi}) \leq \frac{1+|\xi|}{|\xi|}(1+\mathrm{e}^{\gamma}).$$

Furthermore, since $(1+x)/x$ is monotonically decreasing, we have (5.11). This completes the proof. ■

Then Lemma 5.11 is proved as follows.

Proof. From the inequality (2.6) with $i=3$, it follows that

$$\begin{aligned}|F(\zeta)| &\leq \left|\frac{1+\log(1+\mathrm{e}^{\pi\sinh\zeta})}{\log(1+\mathrm{e}^{\pi\sinh\zeta})}\right|^{1-\alpha}\frac{|\psi'_{\mathrm{DE3\ddagger}}(\zeta)|}{|1+\mathrm{e}^{\pi\sinh\zeta}|^{\beta}}\\ &= 2\left|\frac{1+\log(1+\mathrm{e}^{\pi\sinh\zeta})}{\log(1+\mathrm{e}^{\pi\sinh\zeta})}\cdot\frac{1}{1+\mathrm{e}^{-\pi\sinh\zeta}}\right|^{1-\alpha}\frac{K(\pi/2)|\cosh\zeta|}{|1+\mathrm{e}^{-\pi\sinh\zeta}|^{\alpha}|1+\mathrm{e}^{\pi\sinh\zeta}|^{\beta}}.\end{aligned}$$

Then use Lemma 5.12 to obtain the desired result. ■

References

[1] S. Haber, Two formulas for numerical indefinite integration, *Mathematics of Computation*, **60** (1993), 279–296.

[2] M. Muhammad and M. Mori, Double exponential formulas for numerical indefinite integration, *Journal of Computational and Applied Mathematics*, **161** (2003), 431–448.

[3] T. Okayama, T. Matsuo, and M. Sugihara, Error estimates with explicit constants for Sinc approximation, Sinc quadrature and Sinc indefinite integration, *Numerische Mathematik*, (to appear).

[4] F. Stenger, Optimal convergence of minimum norm approximations in H_p, *Numerische Mathematik*, **29** (1978), 345–362.

[5] ———, *Numerical Methods Based on Sinc and Analytic Functions*, Springer-Verlag, New York, 1993.

[6] M. Sugihara, Optimality of the double exponentioal formula—functional analysis approach, *Numerische Mathematik*, **75** (1997), 379–395.

[7] H. Takahasi and M. Mori, Double exponential formulas for numerical integration, *Publications of the Research Institute for Mathematical Sciences,* Kyoto University, **9** (1974), 721–741.

[8] K. Tanaka, M. Sugihara, K. Murota, and M. Mori, Function classes for double exponential integration formulas, *Numerische Mathematik*, **111** (2009), 631–655.

[9] N. Yamanaka, T. Okayama, S. Oishi, and T. Ogita, A fast verified automatic integration algorithm using double exponential formula, *Nonlinear Theory and Its Applications, IEICE*, **1** (2010), 119–132.